

SPACEABILITY OF SETS OF NOWHERE L^q FUNCTIONS

PEDRO L. KAUFMANN AND LEONARDO PELLEGRINI

ABSTRACT. We say that a function $f : [0, 1] \rightarrow \mathbb{R}$ is *nowhere L^q* if, for each nonvoid open subset U of $[0, 1]$, the restriction $f|_U$ is not in $L^q(U)$. For a fixed $1 \leq p < \infty$, we will show that the set

$$S_p \doteq \{f \in L^p[0, 1] : f \text{ is nowhere } L^q, \text{ for each } p < q \leq \infty\},$$

united with $\{0\}$, contains an isometric and complemented copy of ℓ_p . In particular, this improves a result from [2] (which in turn is an improvement of a result from [4]), since S_p turns out to be spaceable. In addition, our result is a generalization of one of the main results from [3].

1. INTRODUCTION

This note is a contribution to the study of large linear structures within essentially non-linear sets of functions which satisfy certain special properties. Given a topological vector space X , it is by now standard to say that a subset $S \subset X$ is *lineable* if $S \cup \{0\}$ contains an infinite dimensional subspace of X , and that S is *spaceable* if $S \cup \{0\}$ contains a *closed* infinite dimensional subspace of X .

Our main object of study will be the $L^p[0, 1]$ spaces. For clearness, we will write L^p instead of $L^p[0, 1]$. Two results motivate the present work. Muñoz-Fernández, Palmberg, Puglisi and Seoane-Sepúlveda showed the following:

Theorem 1.1 (from [4]). *Let $1 \leq p < q$. Then $L^p \setminus L^q$ is $\mathfrak{c}$ -lineable.*

Here, $\mathfrak{c}$ indicates that we can find a vector space with a Hamel basis of cardinality $\mathfrak{c}$ contained in $(L^p \setminus L^q) \cap \{0\}$. The other result is from Botelho, Diniz, Fávoro and Pellegrino:

Theorem 1.2 (from [1]). *For every $p > 0$, $\ell_p \setminus \bigcup_{0 < q < p} \ell_q$ is spaceable.*

In view of Theorems 1.1 and 1.2 it is natural to ask whether we can obtain a result for L^p spaces analogue to Theorem 1.2; Botelho, Fávoro, Pellegrino and Seoane-Sepúlveda have obtained a positive answer to that question:

Theorem 1.3 (from [2]). *For each $p > 0$, $L^p \setminus \bigcup_{q > p} L^q$ is spaceable.*

In this work we will present an improvement of Theorem 1.3, one could say, to the *local* level. We say that $f : [0, 1] \rightarrow \mathbb{R}$ is *nowhere L^q* , for some $1 \leq q \leq \infty$, if for each nonvoid open subset U of $[0, 1]$, $f|_U$ is not in $L^q(U)$. For each $1 \leq p < \infty$, we define

$$(1.1) \quad S_p \doteq \{f \in L^p([0, 1]) : f \text{ is nowhere } L^q, \text{ for each } p < q \leq \infty\};$$

then S_p satisfies the following property, which is our main result:

Theorem 1.4 (Main). *For each $1 \leq p < \infty$, $(S_p \cup \{0\}) \subset L^p$ contains an isometric and complemented copy of ℓ_p . In particular, S_p is spaceable.*

Section 2 will be dedicated to the proof of Theorem 1.4. Note that it only improves Theorem 1.3 for $p \geq 1$. We decided to restrict ourselves to this case for the sake of uniformity of arguments; the careful reader can verify that the same construction can be used to show the spaceability of S_p also for the quasi-Banach $p < 1$ case. Note that the elements from S_p are *nowhere essentially bounded* (that is, nowhere L^∞). It is worth mentioning that Theorem 1.4 generalizes one of the main theorems of [3], which states the following:

2010 *Mathematics Subject Classification.* Primary: 26A30; Secondary: 26A42.

Key words and phrases. Spaceability, nowhere integrable functions.

The first author was supported by CAPES, Research Grant PNPd 2256-2009.

Theorem 1.5 (from [3]). *The set $\mathcal{G}$ of L^1 functions which are nowhere essentially bounded is spaceable in L^1 .*

In the same paper, it was shown that $\mathcal{G}$ is *strongly $\mathfrak{c}$ -algebrable*, that is, $\mathcal{G} \cup \{0\}$ contains a $\mathfrak{c}$ -generated free algebra (with the pointwise multiplication). Let us mention that there are no nontrivial algebraic structures, not even 1-dimensional ones, within S_p , and even within $L^p \setminus L^q$, for $1 \leq p < q < \infty$. In effect, if $f \in L^p \setminus L^q$ for such p and q , then f^n is not in L^p for large enough n .

2. PROOF OF THE MAIN RESULT

We start by discriminating three objects that will be needed in order to construct a nice basic sequence of L^p contained in S_p .

First: an almost disjoint family of Cantor-built sets $(A_n)_n$. First, we recall the construction of the Cantor set of (Lebesgue) measure one half, which we will denote by C . We start with the closed unit interval (denote it by C_1), and remove the center open interval of length $1/4$, obtaining C_2 , a disjoint union of two closed intervals. From each of them, we remove the center open interval of length $1/4^2$, obtaining C_3 , the disjoint union of four closed intervals. Repeating the process and taking the intersection of all C_n , we obtain C . For our convenience, we will call a *hole* of C each open interval that was removed in each step of the construction of C . Let t_n be the right endpoint of the interval which is the one more to the left, among the closed intervals that constitute C_n . The explicit formula for t_n is $\frac{2 + \sum_{j=2}^n 2^{j-2}}{2^{2n-1}}$, and for our purposes what is important about the sequence t_n is the following:

- (1) $t_n < 2^{3-2n}$ and
- (2) $m(C \cap [0, t_n]) = 2^n$.

This information will be needed later on. We now proceed introducing an specific notation for sets which are essentially unions of sets similar to C .

Definition 2.1. (1) For each subinterval I from $[0, 1]$ and each subset $A \subset [0, 1]$, denote

$$A_I \doteq (b - a)A + a;$$

- (2) a nonvoid subset $A \subset [0, 1]$ of the form $\cup\{A_j : j \in \Gamma\}$ is said to be *C-built* (with *C-components* A_j) if there is an almost disjoint family $\{I_j : j \in \Gamma\}$ of subintervals of $[0, 1]$ such that $A_j = C_{I_j}$, where C is the Cantor set of measure $1/2$.

According to the notation from [3], C-built sets are in particular *Cantor-built*, which means, roughly speaking, that they are essentially unions of Cantor sets. The *holes* of a set of the form C_I are defined by the natural way: the set of all holes of C_I is given by $\{J_I : J \text{ is a hole of } C\}$. With the notion of C-built sets in hands we can start partitioning $[0, 1]$. First, put $A_1 \doteq C$. A_2 is defined as follows:

$$A_2 \doteq \cup\{C_J : J \text{ is a hole of } A_1\}.$$

Note that A_1 and A_2 are almost disjoint, and that $m(A_2) = 1/4$. Assuming that we have defined A_n , for some $n \geq 2$, then A_{n+1} is defined inductively:

$$A_{n+1} \doteq \cup\{C_J : J \text{ is a hole of a Cantor component of } A_n\}.$$

This way, we obtain a sequence $(A_n)_n$ of almost disjoint C-built sets satisfying

- (1) $m(A_n) = 2^{-n}$, and
- (2) for any given nonempty open subset U of $[0, 1]$, there exists an $n_0 \in \mathbb{N}$ such that A_n has a C-component contained in U , for each $n \geq n_0$.

Second: a convenient element h of $L^p \setminus \cup_{p < q \leq \infty} L^q$. Let $(r_j)_j$ be a strictly decreasing sequence of real numbers converging to p . For each j , the function $h_j : x \mapsto x^{-1/r_j}$ is in L^p , but not in L^q for any $r_j \leq q \leq \infty$. Define $\tilde{h}$ by

$$(2.1) \quad \tilde{h} \doteq \sum_{j=1}^{\infty} \frac{1}{2^j} \frac{h_j}{\|h_j\|_p}.$$

$\tilde{h}$ is well-defined since the series converges absolutely and L^p is complete. Then $\tilde{h} \in L^p \setminus \cup_{p < q < \infty} L^q$, and moreover:

Lemma 2.2. $h \doteq \chi_C \tilde{h} \in L^p \setminus \cup_{p < q < \infty} L^q$.

Proof. It is easily seen that $h \in L^p$. To show that, for a fixed $p < q < \infty$ we have $h \notin L^q$, consider j_0 such that $r_{j_0} < q$. Then

$$\int_C |h|^q = \int_C |\tilde{h}|^q \geq \int_C \frac{1}{2^{j_0 q}} \frac{h_{j_0}^q}{\|h_{j_0}\|_p^q} \geq \frac{1}{2^{j_0 q} \|h_{j_0}\|_p^q} \int_C x^{-q/r_{j_0}}.$$

We will show that the integral to the right converges to $+\infty$. In effect, denoting $s \doteq q/r_{j_0} > 1$ and considering the sequence t_n defined previously, we have that

$$\begin{aligned} \int_C x^{-q/r_{j_0}} &= \int_C x^{-s} \geq \int_{C \cap [0, t_n]} x^{-s} \geq m(C \cap [0, t_n]) \inf\{x^{-s} : x \in C \cap [0, t_n]\} \\ &\geq m(C \cap [0, t_n]) t_n^{-s} > 2^{-n} 2^{s(2n-3)} > 2^{(2s-1)n}. \end{aligned}$$

Since the right handside tends to infinity along with n , it follows that $h \notin L^q$. The remaining case, $q = \infty$, is automatically covered, since $L^\infty \subset L^q$, for each $q < \infty$. ♠

Third: a disjoint infinite family of strictly increasing sequences of positive integers, $\{(n_j^k)_j : k \in \mathbb{N}\}$. It is a simple task for the reader to convince herself/himself that such family exists.

Before proceeding with the proof of the main result, let us establish the following notation: if $I = [a, b]$ is a subinterval of $[0, 1]$ and $f : [0, 1] \rightarrow \mathbb{R}$, we define $f_I : [0, 1] \rightarrow \mathbb{R}$ by

$$f_I(x) \doteq \begin{cases} f\left(\frac{x-a}{b-a}\right), & \text{if } x \in I; \\ 0, & \text{if } x \notin I. \end{cases}$$

Note that this notation is coherent with Definition 2.1 (1), since for any subset $B \subset [0, 1]$, $(\chi_B)_I = \chi_{B_I}$. It is easily seen that, for any subinterval I from $[0, 1]$ and any $1 \leq p \leq \infty$, f is in L^p if and only if f_I is in L^p . In particular, for each subinterval I from $[0, 1]$, $h_I \in L^p \setminus \cup_{p < q < \infty} L^q$.

Proof (of Theorem 1.4). Consider $(A_j)_j$, h and $\{(n_j^k)_j : k \in \mathbb{N}\}$ as defined above. For a fixed $j \geq 2$, let $(B_l)_l$ be a sequence of all C-components of A_j , and consider their respective convex hulls I_l . Note that the I_l are disjoint. Define

$$f_j \doteq \sum_{l=1}^{\infty} \frac{1}{2^l} \frac{h_{I_l}}{\|h_{I_l}\|_p}.$$

For $j = 1$ just put $f_1 \doteq \frac{h}{\|h\|_p}$. Then each f_j is in L^p with $\|f_j\|_p = 1$, and f_j is not in $L^q(U)$, for any $p < q \leq \infty$ and any open set U containing a C-component of A_j . Note also, by the construction of h , that f_j is zero outside of A_j . For each k , consider

$$g_k \doteq \sum_{j=1}^{\infty} \frac{1}{2^j} f_{n_j^k}.$$

Then $g_k \in S_p$ and $\|g_k\|_p = 1$, and since the functions g_k have almost disjoint supports, it follows that $(g_k)_k$ is a complemented basic sequence in L^p , isometrically equivalent to the canonical basis of ℓ_p . In particular, $\overline{\text{span}(\{g_k : k \in \mathbb{N}\})}$ is complemented in L^p and isometrically isomorphic to ℓ_p .

To complete our proof, it remains to show that $\overline{\text{span}(\{g_k : k \in \mathbb{N}\})} \subset S_p \cup \{0\}$. Any given nonzero element of $\overline{\text{span}(\{g_k : k \in \mathbb{N}\})}$ is of the form

$$g = \sum_{k=1}^{\infty} a_k g_k,$$

where some a_{k_0} is assumed to be nonzero. Since the functions g_k have almost disjoint supports and g_{k_0} is nowhere L^q for any given $p < q \leq \infty$, it follows that $g \in S_p$. ♠

REFERENCES

- [1] G. Botelho, D. Diniz, V. Fávoro, and D. Pellegrino, *Spaceability in Banach and quasi-Banach sequence spaces*, Linear Algebra Appl. **434** (2011), no. 5, 1255–1260.
- [2] G. Botelho, V. Fávoro, D. Pellegrino, and J. B. Seoane-Sepúlveda, *$L_p[0, 1] \setminus \cup_{q>p} L_q[0, 1]$ is spaceable for every $p > 0$* , preprint, 2011.
- [3] S. Głąb, P. L. Kaufmann, and L. Pellegrini, *Spaceability and algebrability of sets of nowhere integrable functions*, preprint, 2011.
- [4] G. A. Muñoz-Fernández, N. Palmberg, D. Puglisi, and J. B. Seoane-Sepúlveda, *Lineability in subsets of measure and function spaces*, Linear Algebra Appl. **428** (2008), 2805–2812.

INSTITUTO DE MATEMÁTICA E ESTATÍSTICA, UNIVERSIDADE DE SÃO PAULO, RUA DO MATÃO, 1010, CEP 05508-900, SÃO PAULO, BRAZIL

E-mail address: `plkaufmann@gmail.com`, `leonardo@ime.usp.br`